arXiv:1704.06997v1 [math.FA] 23 Apr 2017

APPROXIMATION THEOREMS CONNECTED WITH DIFFERENTIAL-DIFFERENCE OPERATOR

CHOKRI ABDELKEFI* AND SAFA CHABCHOUB**

Abstract. In the present paper, we propose to give an extension to the context of Dunkl theory of the notion of translation and in connection with this a corresponding extension of Taylor's formula. More precisely, we prove some properties and estimates of the integral remainder in the generalized Taylor formula associated to the Dunkl operator on the real line and we describe the Besov-type spaces for which the remainder has a given order.

1. Introduction

Delsartes gave in [10, 11] a certain extension of the notion of translation and in connection with this a corresponding extension of Taylor's formula. Generalized translation have later been considered from various points of view by many authors (see Levitan [13], Bochner [8, 9]). Löfström and Peetre in [14] estimated the remainder in the generalized Taylor's formula and they described the space of functions for which the remainder has a given order.

Our aim in this paper is to extend the results obtained in [14] to the context of Dunkl theory. More precisely, we prove some properties and estimates of the integral remainder of order k associated to the Dunkl operator on the real line and we establish the coincidence between two characterizations of Besov-type spaces related to this integral remainder.

For a real parameter $\alpha > -\frac{1}{2}$, the Dunkl operator on the real line denoted by Λ_α, is a differential-difference operator introduced in 1989 by C. Dunkl in [12]. This operator is associated with the reflection group $\mathbb{Z}_2$ on $\mathbb{R}$ and is given by

$$\Lambda_\alpha f(x) = \frac{df}{dx}(x) + \frac{2\alpha+1}{x}\Big[\frac{f(x)-f(-x)}{2}\Big],\ f \in \mathcal{C}^1(\mathbb{R}).$$

The Dunkl operator can be considered as a perturbation of the usual derivative by reflection part. This operator plays a major role in the study of quantum harmonic oscillators governed by Wigner's commutation rules (see [18]). The Dunkl kernel E_α related to Λ_α is used to define the Dunkl transform which enjoys properties similar to those of the classical Fourier transform. The Dunkl kernel E_α satisfies a product formula (see [19]). This allows us to define the Dunkl translation τ_x, $x \in \mathbb{R}$ (see next section). If the parameter $\alpha = -\frac{1}{2}$, then the operator Λ_α reduces to the

1991 *Mathematics Subject Classification.* Primary 44A15, 46E30; Secondary 44A35.

Key words and phrases. Dunkl operator, Dunkl translation operators, Generalized Taylor formula, Besov-Dunkl spaces.

This work was completed with the support of the DGRST research project LR11ES11, University of Tunis El Manar, Tunisia.

differential operator $\frac{df}{dx}$. Therefore Dunkl analysis can be viewed as a generalization of the classical Fourier analysis on $\mathbb{R}$.

In 2003, the classical Taylor formula with integral remainder was extended in [16] to the one dimensional Dunkl operator Λ_α :
For $k = 1, 2, ...,$ $f \in \mathcal{E}(\mathbb{R})$ and $a \in \mathbb{R}$, we have

$$\tau_x(f)(a) = \sum_{p=0}^{k-1} b_p(x)\Lambda_\alpha^p f(a) + R_k(x, f)(a), \quad x \in \mathbb{R}\backslash\{0\},$$

with $R_k(x, f)(a)$ is the integral remainder of order k given by

$$R_k(x, f)(a) = \int_{-|x|}^{|x|} \Theta_{k-1}(x, y)\tau_y(\Lambda_\alpha^k f)(a)A_\alpha(y)dy,$$

where $\mathcal{E}(\mathbb{R})$ is the space of infinitely differentiable functions on $\mathbb{R}$ and $(\Theta_p)_{p\in\mathbb{N}}$, $(b_p)_{p\in\mathbb{N}}$ are two sequences of functions constructed inductively from the function A_α defined on $\mathbb{R}$ by $A_\alpha(x) = |x|^{2\alpha+1}$ (see next section).

There are many ways to define the Besov spaces (see [6, 7, 17]) and the Besov-Dunkl spaces (see [1, 2, 3, 4]). It is well known that Besov spaces can be described by means of differences using the modulus of continuity of functions. These spaces defined by the modulus of smoothness occur more naturally in many areas of analysis including approximation theory.

In this paper we define the following weighted function spaces:
Let $0 < \beta < 1$, $1 \le p < +\infty$, $1 \le q \le +\infty$ and k a positive integer ($k = 1, 2, ...$).
• We denote by $L^p(\mu_\alpha)$ the space of complex-valued functions f, measurable on $\mathbb{R}$ such that

$$\|f\|_{p,\alpha} = \left(\int_{\mathbb{R}} |f(x)|^p d\mu_\alpha(x)\right)^{1/p} < +\infty,$$

where μ_α is a weighted Lebesgue measure associated to the Dunkl operator given by

$$d\mu_\alpha(x) = \frac{A_\alpha(x)}{2^{\alpha+1}\Gamma(\alpha+1)}dx,$$

with A_α is the function defined on $\mathbb{R}$ by

$$A_\alpha(x) = |x|^{2\alpha+1}, \quad x \in \mathbb{R}.$$

• The Besov-Dunkl space of order k denoted by $\mathcal{B}^k\mathcal{D}_{p,q}^{\beta,\alpha}$ is the subspace of functions f in $\mathcal{E}(\mathbb{R}) \cap L^p(\mu_\alpha)$ such that $\Lambda_\alpha^{k-1} f \in L^p(\mu_\alpha)$ and satisfying

$$\int_0^{+\infty} \Big(\frac{\omega_{p,\alpha}^k(x, f)}{x^{\beta+k-1}}\Big)^q \frac{dx}{x} < +\infty \quad if \quad q < +\infty$$

$$\text{and} \qquad \sup_{x>0} \frac{\omega_{p,\alpha}^k(x, f)}{x^{\beta+k-1}} < +\infty \quad if \quad q = +\infty,$$

with $\omega_{p,\alpha}^k(x, f) = \sup_{|y|\le x} \|R_{k-1}(y, f) - b_{k-1}(y)\Lambda_\alpha^{k-1} f\|_{p,\alpha}$, where we put $\Lambda_\alpha^0 f = f$ and $R_0(x, f) = \tau_x(f)$, for $k = 1$.
• Put $\mathcal{D}_{p,\alpha}^k$ the subspace of functions f in $\mathcal{E}(\mathbb{R}) \cap L^p(\mu_\alpha)$ such that $\Lambda_\alpha^k f$ are in

$L^p(\mu_\alpha)$. We consider the subspace $\mathcal{K}^k\mathcal{D}^{\beta,\alpha}_{p,q}$ of functions $f \in \mathcal{D}^{k-1}_{p,\alpha} + \mathcal{D}^k_{p,\alpha}$ satisfying

$$\int_0^{+\infty} \Big(\frac{K^k_{p,\alpha}(x,f)}{x^\beta}\Big)^q \frac{dx}{x} < +\infty \quad if \quad q < +\infty$$

$$\text{and} \qquad \sup_{x>0} \frac{K^k_{p,\alpha}(x,f)}{x^\beta} < +\infty \quad if \quad q = +\infty,$$

where $K^k_{p,\alpha}$ is the Peetre K-functional given by

$$K^k_{p,\alpha}(x,f) = \inf_{f=f_0+f_1} \Big\{ \|\Lambda^{k-1}_\alpha f_0\|_{p,\alpha} + x\|\Lambda^k_\alpha f_1\|_{p,\alpha},\ f_0 \in \mathcal{D}^{k-1}_{p,\alpha},\ f_1 \in \mathcal{D}^k_{p,\alpha} \Big\}.$$

The contents of the present paper are as follows.
In section 2, we collect some basic definitions and results about harmonic analysis associated with the Dunkl operator Λ_α.
In section 3, we give some properties and estimates of the integral remainder of order k. Finally, we establish that

$$\mathcal{B}^k\mathcal{D}^{\beta,\alpha}_{p,q} = \mathcal{K}^k\mathcal{D}^{\beta,\alpha}_{p,q}.$$

Along this paper, we use c to represent a suitable positive constant which is not necessarily the same in each occurrence.

2. Preliminaries

In this section, we recall some notations and results in Dunkl theory on $\mathbb{R}$ and we refer for more details to [5, 12, 19].

For $\lambda \in \mathbb{C}$, the initial problem

$$\Lambda_\alpha(f)(x) = \lambda f(x), \quad f(0) = 1, \quad x \in \mathbb{R},$$

has a unique solution $E_\alpha(\lambda.)$ called Dunkl kernel given by

$$E_\alpha(\lambda x) = j_\alpha(i\lambda x) + \frac{\lambda x}{2(\alpha+1)} j_{\alpha+1}(i\lambda x), \quad x \in \mathbb{R},$$

where j_α is the normalized Bessel function of the first kind and order α.

The Dunkl kernel E_α satisfies the following product formula

$$E_\alpha(ixt)E_\alpha(iyt) = \int_{\mathbb{R}} E_\alpha(itz)d\gamma_{x,y}(z), \quad x,y,t \in \mathbb{R},$$

where $\gamma_{x,y}$ is a signed measure on $\mathbb{R}$ with compact support.

For $x, y \in \mathbb{R}$ and f a continuous function on $\mathbb{R}$, the Dunkl translation operator τ_x is given by

$$\tau_x(f)(y) = \int_{\mathbb{R}} f(z)d\gamma_{x,y}(z).$$

.
The Dunkl translation operator satisfies the following properties :

(1) τ_x is a continuous linear operator from $\mathcal{E}(\mathbb{R})$ into itself.
(2) For all $f \in \mathcal{E}(\mathbb{R})$, we have

$$\tau_x(f)(y) = \tau_y(f)(x) \quad \text{and} \quad \tau_0(f)(x) = f(x)$$

$$\tau_x \, o \, \tau_y = \tau_y \, o \, \tau_x \quad \text{and} \quad \Lambda_\alpha \, o \, \tau_x = \tau_x \, o \, \Lambda_\alpha. \tag{2.1}$$

(3) For all $x \in \mathbb{R}$, the operator τ_x extends to $L^p(\mu_\alpha)$, $p \geq 1$ and we have for f in $L^p(\mu_\alpha)$

$$\|\tau_x(f)\|_{p,\alpha} \leq \sqrt{2}\|f\|_{p,\alpha}. \tag{2.2}$$

It has been shown in [16], the following generalized Taylor formula with integral remainder:

Proposition 2.1. *For $k = 1, 2, ...$, $f \in \mathcal{E}(\mathbb{R})$ and $a \in \mathbb{R}$, we have*

$$\tau_x f(a) = \sum_{p=0}^{k-1} b_p(x)\Lambda_\alpha^p f(a) + R_k(x, f)(a), \quad x \in \mathbb{R}\backslash\{0\}, \tag{2.3}$$

with $R_k(x, f)(a)$ is the integral remainder of order k given by

$$R_k(x, f)(a) = \int_{-|x|}^{|x|} \Theta_{k-1}(x, y)\tau_y(\Lambda_\alpha^k f)(a)A_\alpha(y)dy, \tag{2.4}$$

where

(i) $b_{2m}(x) = \dfrac{1}{(\alpha+1)_m m!}\Big(\dfrac{x}{2}\Big)^{2m}$ *and* $b_{2m+1}(x) = \dfrac{1}{(\alpha+1)_{m+1} m!}\Big(\dfrac{x}{2}\Big)^{2m+1}$, *for* $m \in \mathbb{N}$.

(ii) $\Theta_{k-1}(x, y) = u_{k-1}(x, y) + v_{k-1}(x, y)$ *with* $u_0(x, y) = \dfrac{sgn(x)}{2A_\alpha(x)}$,

$v_0(x, y) = \dfrac{sgn(y)}{2A_\alpha(y)}$, $u_k(x, y) = \displaystyle\int_{|y|}^{|x|} v_{k-1}(x, z)dz$ *and*

$v_k(x, y) = \dfrac{sgn(y)}{A_\alpha(y)} \displaystyle\int_{|y|}^{|x|} u_{k-1}(x, z)A_\alpha(z)dz.$

According to ([20], Lemma 2.2), the Dunkl operator Λ_α have the following regularity properties:

$$\Lambda_\alpha \text{ leaves } \mathcal{C}_c^\infty(\mathbb{R}) \text{ and the Schwartz space } \mathcal{S}(\mathbb{R}) \text{ invariant.} \tag{2.5}$$

3. Characterizations of Besov-Dunkl spaces of order k

In this section, we start with the proof of some properties and estimates of the integral remainder in the generalized Taylor formula.

Remark 3.1. *Let $k = 1, 2, ...$, $f \in \mathcal{E}(\mathbb{R})$ and $x \in \mathbb{R}\backslash\{0\}$.*

(1) From Proposition 2.1, we have

$$\begin{aligned} R_k(x, f) &= \tau_x(f) - f - b_1(x)\Lambda_\alpha f ... - b_{k-1}(x)\Lambda_\alpha^{k-1} f \\ &= R_{k-1}(x, f) - b_{k-1}(x)\Lambda_\alpha^{k-1} f, \end{aligned} \tag{3.1}$$

where we put for $k = 1$, $R_0(x, f) = \tau_x(f)$. Observe that

$$R_1(x, f) = R_0(x, f) - b_0(x)\Lambda_\alpha^0 f = \tau_x(f) - f.$$

(2) According to ([16], p.352) and Proposition 2.1, (i), we have

$$\begin{aligned} \int_{-|x|}^{|x|} |\Theta_{k-1}(x, y)|A_\alpha(y)dy &\leq b_k(|x|) + |x|b_{k-1}(|x|) \\ &\leq c\,|x|^k. \end{aligned} \tag{3.2}$$

(3) Note that the function $y \longmapsto \tau_y(f) - f$ is continuous on $\mathbb{R}$ (see [15], Lemma 1, (ii)), which implies that the same is true for the function $y \longmapsto R_k(y, f)$.

Lemma 3.1. *Let $k = 1, 2, ...$, then there exists a constant $c > 0$ such that for all $f \in \mathcal{E}(\mathbb{R})$ satisfying $\Lambda_\alpha^{k-1} f \in L^p(\mu_\alpha)$, we have*

$$\|R_{k-1}(x, f)\|_{p,\alpha} \leq c\,|x|^{k-1} \|\Lambda_\alpha^{k-1} f\|_{p,\alpha}, \quad x \in \mathbb{R}\backslash\{0\}. \tag{3.3}$$

Proof. Let $k = 1, 2, ...$, $f \in \mathcal{E}(\mathbb{R})$ such that $\Lambda_\alpha^{k-1} f \in L^p(\mu_\alpha)$ and $x \in \mathbb{R}\backslash\{0\}$. For $k = 1$, by (2.2), it's clear that $\|R_0(x, f)\| = \|\tau_x(f)\|_{p,\alpha} \leq c\|f\|_{p,\alpha}$. Using Minkowski's inequality for integrals, (2.2) and (2.4), we have for $k \geq 2$

$$\begin{aligned}\|R_{k-1}(x, f)\|_{p,\alpha} &\leq \int_{-|x|}^{|x|} |\Theta_{k-2}(x, y)|\ \|\tau_y(\Lambda_\alpha^{k-1} f)\|_{p,\alpha} A_\alpha(y)dy \\ &\leq c\ \|\Lambda_\alpha^{k-1} f\|_{p,\alpha} \int_{-|x|}^{|x|} |\Theta_{k-2}(x, y)| A_\alpha(y)dy.\end{aligned}$$

From (3.2), we deduce our result. □

Remark 3.2. *Let $k = 1, 2, ...$, $f \in \mathcal{E}(\mathbb{R})$ such that $\Lambda_\alpha^{k-1} f \in L^p(\mu_\alpha)$. Then for $x \in \mathbb{R}\backslash\{0\}$, we have by (3.1), (3.3) and Proposition 2.1,*

$$\begin{aligned}\|R_k(x, f)\|_{p,\alpha} &\leq \|R_{k-1}(x, f)\|_{p,\alpha} + \|b_{k-1}(x)\Lambda_\alpha^{k-1} f\|_{p,\alpha} \\ &\leq c\,|x|^{k-1} \|\Lambda_\alpha^{k-1} f\|_{p,\alpha}.\end{aligned} \tag{3.4}$$

Lemma 3.2. *For $x \in \mathbb{R}\backslash\{0\}$ and $p \in \mathbb{N}$, we have*

$$\int_{-|x|}^{|x|} \Theta_0(x, y) b_p(y) A_\alpha(y)dy = b_{p+1}(x). \tag{3.5}$$

Proof. Let $x \in \mathbb{R}\backslash\{0\}$. Using Proposition 2.1, we have:
• If $p = 2m$, $m \in \mathbb{N}$,

$$\begin{aligned}&\int_{-|x|}^{|x|} \Theta_0(x, y) b_{2m}(y) A_\alpha(y)dy \\ &= \int_{-|x|}^{|x|} \frac{sgn(x)|y|^{2\alpha+1}}{2|x|^{2\alpha+1}} b_{2m}(y)dy + \int_{-|x|}^{|x|} \frac{sgn(y)}{2} b_{2m}(y)dy \\ &= \frac{x}{2^{2m}|x|^{2\alpha+2}(\alpha+1)_m m!} \int_0^{|x|} y^{2\alpha+2m+1} dy \\ &= \frac{x}{2^{2m}(\alpha+1)_m m!} \frac{|x|^{2m}}{2(\alpha+m+1)} \\ &= b_{2m+1}(x).\end{aligned}$$

• If $p = 2m + 1$, $m \in \mathbb{N}$, we get

$$\begin{aligned}
&\int_{-|x|}^{|x|} \Theta_0(x,y) b_{2m+1}(y) A_\alpha(y) dy \\
&\quad = \int_{-|x|}^{|x|} \frac{sgn(x)|y|^{2\alpha+1}}{2|x|^{2\alpha+1}} b_{2m+1}(y) dy + \int_{-|x|}^{|x|} \frac{sgn(y)}{2} b_{2m+1}(y) dy \\
&\quad = \frac{1}{2^{2m+1}(\alpha+1)_{m+1} m!} \int_0^{|x|} y^{2m+1} dy \\
&\quad = \frac{1}{2^{2m+1}(\alpha+1)_{m+1} m!} \frac{|x|^{2m+2}}{2(m+1)} \\
&\quad = b_{2m+2}(x).
\end{aligned}$$

Hence the Lemma is proved. □

Lemma 3.3. *Let $k = 1, 2, ...,$ $f \in \mathcal{E}(\mathbb{R})$, $x \in \mathbb{R}\backslash\{0\}$ and $a \in \mathbb{R}$. Then we have,*

$$R_k(x,f)(a) = \int_{-|x|}^{|x|} \Theta_0(x,y) R_{k-1}(y, \Lambda_\alpha f)(a) A_\alpha(y) dy. \tag{3.6}$$

Proof. Let $k = 1, 2, ...,$ $f \in \mathcal{E}(\mathbb{R})$, $x \in \mathbb{R}\backslash\{0\}$ and $a \in \mathbb{R}$. We have from (2.3), (2.4) and the fact that $R_0(y, \Lambda_\alpha f)(a) = \tau_y(\Lambda_\alpha f)$,

$$R_1(x,f)(a) = (\tau_x(f) - f)(a) = \int_{-|x|}^{|x|} \Theta_0(x,y) \tau_y(\Lambda_\alpha f)(a) A_\alpha(y) dy,$$

hence the property (3.6) is true for $k = 1$.
Suppose that

$$R_k(x,f)(a) = \int_{-|x|}^{|x|} \Theta_0(x,y) R_{k-1}(y, \Lambda_\alpha f)(a) A_\alpha(y) dy,$$

then by (3.1) and (3.5), we get

$$\begin{aligned}
&\int_{-|x|}^{|x|} \Theta_0(x,y) R_k(y, \Lambda_\alpha f)(a) A_\alpha(y) dy \\
&\quad = \int_{-|x|}^{|x|} \Theta_0(x,y) R_{k-1}(y, \Lambda_\alpha f)(a) A_\alpha(y) dy \\
&\quad - \int_{-|x|}^{|x|} \Theta_0(x,y) b_{k-1}(y) \Lambda_\alpha^k f(a) A_\alpha(y) dy \\
&\quad = R_k(x,f)(a) - \Lambda_\alpha^k f(a) \int_{-|x|}^{|x|} \Theta_0(x,y) b_{k-1}(y) A_\alpha(y) dy \\
&\quad = R_k(x,f)(a) - b_k(x) \Lambda_\alpha^k f(a) \\
&\quad = R_{k+1}(x,f)(a).
\end{aligned}$$

By induction, we deduce our result. □

Lemma 3.4. *Let $k = 1, 2, ...,$ $f \in \mathcal{E}(\mathbb{R})$, $x \in \mathbb{R}\backslash\{0\}$ and $a \in \mathbb{R}$. We denote by*

$$I_1(x,f)(a) = \int_{-|x|}^{|x|} \Theta_0(x,y) \tau_y(f)(a) A_\alpha(y) dy,$$

and for $k \geq 2$

$$I_k(x,f)(a) = \int_{-|x|}^{|x|} \Theta_0(x,y) I_{k-1}(y,f)(a) A_\alpha(y) dy.$$

Then, we have

$$\Lambda_\alpha^{k+1}\big(I_k(x,f)\big)(a) = \Lambda_\alpha^k\big(I_k(x,\Lambda_\alpha f)\big)(a) \tag{3.7}$$

$$\text{and} \qquad \Lambda_\alpha^k I_k(x,f)(a) = R_k(x,f)(a). \tag{3.8}$$

Proof. Let $k = 1, 2, ...,$ $f \in \mathcal{E}(\mathbb{R})$, $x \in \mathbb{R}\backslash\{0\}$ and $a \in \mathbb{R}$.
• Using (2.1), we have

$$\begin{aligned}\Lambda_\alpha^2\big(I_1(x,f)\big)(a) &= \int_{-|x|}^{|x|} \Theta_0(x,y)\Lambda_\alpha\tau_y(\Lambda_\alpha f)(a)A_\alpha(y)dy \\ &= \Lambda_\alpha\big(I_1(x,\Lambda_\alpha f)\big)(a).\end{aligned}$$

Suppose that

$$\Lambda_\alpha^{k+1}\big(I_k(x,f)\big)(a) = \Lambda_\alpha^k\big(I_k(x,\Lambda_\alpha f)\big)(a),$$

this gives

$$\begin{aligned}\Lambda_\alpha^{k+2}\big(I_{k+1}(x,f)\big)(a) &= \int_{-|x|}^{|x|} \Theta_0(x,y)\Lambda_\alpha\big(\Lambda_\alpha^{k+1} I_k(y,f)\big)(a)A_\alpha(y)dy \\ &= \int_{-|x|}^{|x|} \Theta_0(x,y)\Lambda_\alpha\big(\Lambda_\alpha^{k} I_k(y,\Lambda_\alpha f)\big)(a)A_\alpha(y)dy \\ &= \Lambda_\alpha^{k+1}\big(I_{k+1}(x,\Lambda_\alpha f)\big)(a).\end{aligned}$$

Then, we obtain (3.7) by induction.
• From (2.1) and (2.4), we can write

$$\begin{aligned}\Lambda_\alpha\big(I_1(x,f)\big)(a) &= \int_{-|x|}^{|x|} \Theta_0(x,y)\Lambda_\alpha(\tau_y f)(a)A_\alpha(y)dy \\ &= \int_{-|x|}^{|x|} \Theta_0(x,y)\tau_y(\Lambda_\alpha f)(a)A_\alpha(y)dy \\ &= R_1(x,f)(a).\end{aligned}$$

Suppose that

$$\Lambda_\alpha^k\big(I_k(x,f)\big)(a) = R_k(x,f)(a),$$

then by (3.6) and (3.7), we have

$$\begin{aligned}\Lambda_\alpha^{k+1}\big(I_{k+1}(x,f)\big)(a) &= \int_{-|x|}^{|x|} \Theta_0(x,y)\Lambda_\alpha^{k+1}\big(I_k(y,f)\big)(a)A_\alpha(y)dy \\ &= \int_{-|x|}^{|x|} \Theta_0(x,y)\Lambda_\alpha^{k}\big(I_k(y,\Lambda_\alpha f)\big)(a)A_\alpha(y)dy \\ &= \int_{-|x|}^{|x|} \Theta_0(x,y)R_k(y,\Lambda_\alpha f)(a)A_\alpha(y)dy \\ &= R_{k+1}(x,f)(a).\end{aligned}$$

By induction, this gives (3.8). □

Before establishing that $\mathcal{B}^k\mathcal{D}^{\beta,\alpha}_{p,q} = \mathcal{K}^k\mathcal{D}^{\beta,\alpha}_{p,q}$, we give a remark, a proposition containing sufficient conditions and an example.

Remark 3.3. *For $k = 1, 2, ...$, $f \in \mathcal{E}(\mathbb{R}) \cap L^p(\mu_\alpha)$ such that $\Lambda^{k-1}_\alpha f$ is in $L^p(\mu_\alpha)$ and $x \in (0, +\infty)$, we can assert from (3.1) that*

(1) $\omega^k_{p,\alpha}(x, f) = \sup\limits_{|y|\le x} \|R_k(y, f)\|_{p,\alpha}$.

(2) For $k = 1$, we have $\omega^k_{p,\alpha}(x, f) = \sup\limits_{|y|\le x} \|\tau_y(f) - f\|_{p,\alpha}$.

Proposition 3.1. *Let $k = 1, 2, ...$, $1 \le p < +\infty$, $1 \le q \le +\infty$, $0 < \beta < 1$ and $f \in \mathcal{E}(\mathbb{R}) \cap L^p(\mu_\alpha)$. If $\Lambda^{k-1}_\alpha f$ and $\Lambda^k_\alpha f$ are in $L^p(\mu_\alpha)$, then $f \in \mathcal{B}^k\mathcal{D}^{\beta,\alpha}_{p,q}$.*

Proof. Let $k = 1, 2, ...$, $1 \le p < +\infty$, $1 \le q \le +\infty$, $0 < \beta < 1$ and $f \in \mathcal{E}(\mathbb{R})$ such that $\Lambda^{k-1}_\alpha f$, $\Lambda^k_\alpha f$ are in $L^p(\mu_\alpha)$. For $x \in (0, +\infty)$, we obtain by (3.3) and (3.4),

$$\omega^k_{p,\alpha}(x, f) \le c\, x^k \|\Lambda^k_\alpha f\|_{p,\alpha} \quad \text{and} \quad \omega^k_{p,\alpha}(x, f) \le c\, x^{k-1} \|\Lambda^{k-1}_\alpha f\|_{p,\alpha}.$$

Then we can write,

$$\int_0^{+\infty} \Big(\frac{\omega^k_{p,\alpha}(x, f)}{x^{\beta+k-1}}\Big)^q \frac{dx}{x} \le c \int_0^1 \Big(\frac{\|\Lambda^k_\alpha f\|_{p,\alpha}}{x^{\beta-1}}\Big)^q \frac{dx}{x} + c \int_1^{+\infty} \Big(\frac{\|\Lambda^{k-1}_\alpha f\|_{p,\alpha}}{x^{\beta}}\Big)^q \frac{dx}{x},$$

giving two finite integrals. Here when $q = +\infty$, we make the usual modification. □

Example 3.1. *From (2.5) and Proposition 3.1, we can assert that the spaces $\mathcal{C}^\infty_c(\mathbb{R})$ and $\mathcal{S}(\mathbb{R})$ are included in $\mathcal{B}^k\mathcal{D}^{\beta,\alpha}_{p,q}$.*

Theorem 3.1. *Let $0 < \beta < 1$, $k = 1, 2, ...$, $1 \le p < +\infty$ and $1 \le q \le +\infty$, then*

$$\mathcal{B}^k\mathcal{D}^{\beta,\alpha}_{p,q} = \mathcal{K}^k\mathcal{D}^{\beta,\alpha}_{p,q}.$$

Proof. We start with the proof of the inclusion $\mathcal{K}^k\mathcal{D}^{\beta,\alpha}_{p,q} \subset \mathcal{B}^k\mathcal{D}^{\beta,\alpha}_{p,q}$. Let f a function in $\mathcal{K}^k\mathcal{D}^{\beta,\alpha}_{p,q}$. If $f = f_0 + f_1$, $f_0 \in \mathcal{D}^{k-1}_{p,\alpha}$ and $f_1 \in \mathcal{D}^k_{p,\alpha}$ is any decomposition of f, we have by (3.3)

$$\begin{aligned} \omega^k_{p,\alpha}(x, f_1) &= \sup_{|y|\le x} \|R_k(y, f_1)\|_{p,\alpha} \\ &\le c\; x^k \|\Lambda^k_\alpha f_1\|_{p,\alpha}, \quad x \in (0, +\infty). \end{aligned} \tag{3.9}$$

Using (3.4), we obtain

$$\begin{aligned} \omega^k_{p,\alpha}(x, f_0) &\le \sup_{|y|\le x} \|R_{k-1}(y, f_0)\|_{p,\alpha} + \sup_{|y|\le x} \|b_{k-1}(y)\Lambda^{k-1}_\alpha f_0\|_{p,\alpha} \\ &\le c\; x^{k-1} \|\Lambda^{k-1}_\alpha f_0\|_{p,\alpha}, \quad x \in (0, +\infty). \end{aligned} \tag{3.10}$$

Hence by (3.9) et (3.10), we deduce that

$$\omega^k_{p,\alpha}(x, f) \le c\; x^{k-1} K^k_{p,\alpha}(x, f),$$

then, $f \in \mathcal{B}^k\mathcal{D}^{\beta,\alpha}_{p,q}$.

To prove the inclusion $\mathcal{B}^k\mathcal{D}^{\beta,\alpha}_{p,q} \subset \mathcal{K}^k\mathcal{D}^{\beta,\alpha}_{p,q}$, we have to make for $f \in \mathcal{B}^k\mathcal{D}^{\beta,\alpha}_{p,q}$ a proper choice of f_0 and f_1. We take for $x \in (0, +\infty)$

$$f_1 = \frac{1}{b_k(x)}\; I_k(x, f).$$

Using (3.8), we obtain

$$x\|\Lambda_\alpha^k f_1\|_{p,\alpha} \leq \; x\,\big(b_k(x)\big)^{-1}\omega_{p,\alpha}^k(x,f) \leq c\,\frac{\omega_{p,\alpha}^k(x,f)}{x^{k-1}}. \tag{3.11}$$

On the other hand, put $f_0 = f - f_1$, we can write by (3.5)

$$f_0 = -\frac{1}{b_k(x)}\int_{-x}^{x}\Theta_0(x,y)\big(I_{k-1}(y,f) - b_{k-1}(y)f\big)A_\alpha(y)dy.$$

From (3.1) and (3.8), we obtain

$$\Lambda_\alpha^{k-1} f_0 = -\frac{1}{b_k(x)}\int_{-x}^{x}\Theta_0(x,y)R_k(y,f)A_\alpha(y)dy.$$

By Minkowski's inequality for integrals and (3.2), we get

$$\begin{aligned}\|\Lambda_\alpha^{k-1} f_0\|_{p,\alpha} &\leq \big(b_k(x)\big)^{-1}\int_{-x}^{x}|\Theta_0(x,y)|\;\|R_k(y,f)\|_{p,\alpha}A_\alpha(y)dy \\ &\leq c\,x^{-k}\omega_{p,\alpha}^k(x,f)\int_{-x}^{x}|\Theta_0(x,y)|\;A_\alpha(y)dy \\ &\leq c\,\frac{\omega_{p,\alpha}^k(x,f)}{x^{k-1}}.\end{aligned} \tag{3.12}$$

By (3.11) et (3.12), we deduce that

$$K_{p,\alpha}^k(x,f) \leq c\,\frac{\omega_{p,\alpha}^k(x,f)}{x^{k-1}},$$

then, $f \in \mathcal{K}^k\mathcal{D}_{p,q}^{\beta,\alpha}$ which completes the proof of the theorem. □

References

[1] C. Abdelkefi and M. Sifi, Characterization of Besov spaces for the Dunkl operator on the real line. Journal of Inequalities in Pure and Applied Mathematics, JIPAM, Vol. 8, Issue 3, 11 pp (2007).

[2] C. Abdelkefi, Dunkl transform on Besov spaces and Herz spaces. Communication in Mathematical Analysis, Vol. 2, No 2, 35-41 (2007).

[3] C. Abdelkefi, J. Ph. Anker, F. Sassi and M. Sifi, Besov-type spaces on $\mathbb{R}^d$ and integrability for the Dunkl transform. Symmetry, Integrability and Geometry: Methods and Applications, SIGMA 5, 019, 15 pages (2009).

[4] C. Abdelkefi, Weighted function spaces and Dunkl transform. Mediterr. J. Math. 9, 499-513 Springer (2012).

[5] B. Amri, J. Ph. Anker, and M. Sifi, Three results in Dunkl analysis. Colloq. Math. 118, 1, 299-312 (2010).

[6] J. L. Ansorena and O. Blasco, Characterization of weighted Besov spaces. Math. Nachr. 171, 5-17 (1995).

[7] O. V. Besov, On a family of function spaces in connection with embeddings and extentions. Trudy Mat. Inst. Steklov 60, 42-81 (1961).

[8] S. Bochner, Positive zonal functions on spheres. Proc. Nat. Acad. Sci., Washihgton, 40, 1141-1147 (1954).

[9] S. Bochner, Sturm-Liouville and Heat equations whose eigenfunctions are ultraspherical polynomials or associated Bessel functions. Proc. Conf. Diff. Eq., Univ. of Maryland, (1955).

[10] J. Delsarte, Sur une extention de la formule de Taylor. J. Math. Pures Appl. 17, 213-231 (1936).

[11] J. Delsarte, Une extention nouvelle de la thorie des fonctions priodiques de Bohr. Acta Math. 69, 259-317 (1938).

[12] C.F. Dunkl, Differential-difference operators associated to reflexion groups. Trans. Amer. Math. Soc., 311, 167-183 (1989).
[13] B. M. Levitan, Generalized translation operators with applications. (In russian) Moscow: Gosudarstvennoe Izdatel'stvo Fiziko-Matematicheskoi Literatury (1962).
[14] J. Löfström, J. Peetre, Approximation theorems connected with generalized translations. Math. Ann., 181, 255-268 (1969).
[15] M.A. Mourou and K. Trimèche, Calderon's reproducing formula related to the Dunkl operator on the real line. Monatshefte für Mathematik 136, 47-65 (2002).
[16] M.A. Mourou, Taylor series associated with a differential-difference operator on the real line. J. Comp. and Appl. Math., 153, 343-354 (2003).
[17] J. Peetre, New thoughts on Besov spaces. Duke Univ. Math. Series, Durham, NC, (1976).
[18] M. Rosenblum, Generalized Hermite polynomials and the Bose-like oscillator calculus. Operator Theory: Advances and Applications, Birkhäuser Verlag, Basel, p. 369-396 (1994).
[19] M. Rösler, Bessel-Type signed hypergroup on $\mathbb{R}$, in Probability measure on groups and related structures. Proc. Conf. Oberwolfach, (1994), H. Heyer and A. Mukherjea (Eds.), World scientific Publ., pp. 292-304 (1995).
[20] M. Rösler, Dunkl operators: theory and applications, in Orthogonal polynomials and special functions (Leuven, 2002). Lect. Notes Math. 1817, Springer-Verlag, 93-135 (2003).

* Department of Mathematics
Preparatory Institute of Engineer Studies of Tunis
1089 Monfleury Tunis, University of Tunis
Tunisia
E-mail address: chokri.abdelkefi@yahoo.fr

** Department of Mathematics
Faculty of Sciences of Tunis
1060 Tunis, University of Tunis El Manar
Tunisia
E-mail address: safachabchoub@yahoo.com